

Authors' manuscript

Booth, D.A., & Knibb, R.C. (2000). What's the evidence that your problem comes from that food? *Food Allergy & Intolerance* 1, 191-200. ([Questionnaire in Appendix](#))

What's the evidence that your problem comes from that food?

David A. Booth* and Rebecca C. Knibb

*Nutritional Psychology Research Group, Psychology-in-Medicine Research Consortium,
The University of Birmingham, Edgbaston, Birmingham B15 2TT, UK*

**Corresponding Author*

The recall of personal experiences relevant to a claim of food allergy or sensitivity is assessed by a psychologically validated tool for evidence that the suspected food could have caused the adverse symptom suffered. If the memory is of actual events that do point to the food as a potential cause of the symptom and the symptom is sufficiently serious, the tool user is recommended to seek testing at a hospital clinic by an immunologist or by the appropriate specialist for a non-allergic sensitivity. If what is recalled does not support the logical possibility of a causal connection between eating that food and occurrence of the symptom, then the user of the tool is pointed to other potential sources of the problem and to other remedies than avoidance of the food that had been blamed.

Unpleasant symptoms are blamed on foods by a substantial minority of European and North American populations. Yet objective tests confirm only a small proportion of these complaints (1).

A total of 300 complainants of food-induced "allergy or sensitivity" were recruited from the community by randomised mailings to electors. Most of the recruits can be assumed (1) to have been in error in attributing their nasty reaction to the food or drink that they named. Results showed that these misattributed symptoms were not usually "psychological", as widely stated in this field, i.e. "an unpleasant reaction to food which is caused by emotions associated with the food rather than by the food itself" (2). Follow-up questionnaires to these recruits and to other research volunteers who did not blame foods for their illnesses detected very little more anxiety or depression among those who supposed that they were food sensitive or allergic than among those controls (3). Rather, "false food allergy" in the community is real suffering with a physical basis (whether or not mediated by the immune system) and, although the cause often is not the food(s) supposed, there was nothing unreasonable in the suspicions elucidated by detailed responses to structured interviews.

Many of the interviewees had indeed been advised by their doctor to look at their diet for a possible cause of their symptoms. Health pages, leaflets and broadcasts of higher or lower reliability alert the general public to the serious reactions that some children or adults have to a great variety of foods and drinks. There are books for the lay reader that go through the scientific evidence for adverse effects from common dietary constituents (e.g. 4).

None of this, however, provides individuals with a basis for judging whether or not a symptom that can come from eating a food, however rarely, has or has not in fact been caused by that food in their cases. There are so many possibilities in the diet, and of other influences on such a symptom, that even an appropriately specialised practitioner often cannot make a reliable diagnosis without blind physical testing. Eating and drinking are so frequent and involve so many combinations of chemically complicated materials that some dietary constituent or another is liable to be consumed before the onset or exacerbation of an illness from any cause. Thus even a highly intelligent, very careful and well informed member of the public is liable to make the mistake of suspecting an innocent food of generating the adverse

symptom. The imputation of hypochondria and worse to millions of ordinary people must therefore be deplored.

Indeed, the great prevalence of misperceived food allergy or sensitivity illustrates the fact evident in other areas of public health (5,6) that the communication of nutrition information on the basis only of general biochemical and epidemiological evidence is not only of little use but can be counterproductive. Advice on healthy eating patterns and other complex choices must be personally tailored from direct evidence of differences between individuals in the outcomes of behaviour. Psychological science is required as well as medical science.

Cognitive psychologists' research findings on memory were applied to interviewees' attempts to recall personal experiences of the symptom in question occurring shortly after consumption of the blamed food (7). Memory for an actual occasion when eating the food was followed by the symptom could be distinguished from the belief in an allergy or sensitivity based only on the theoretical knowledge that such a food can give rise to the symptom in question.

That set of cognitive performance criteria for the actuality of a food-symptom episode was supplemented by criteria based on the logic of evidence for causal connections of any type (7,8). These causal criteria are of two sorts. Causal contingency requires that the recalled co-occurrence of food and symptom was unlikely to be mere coincidence arising from frequent occurrence of either the food or the symptom. The contiguity requirement is that the symptom arose sufficiently soon after the food had been eaten for some sort of cause-effect relationship to be plausible. The criteria of contiguity were however applied to the interview data without reference to current knowledge of pathophysiological mechanisms specific to that food and symptom (7).

In nearly all cases, interviewees whose food-symptom recall met most of the criteria of actuality, contingency or contiguity could be assigned to the same category by just one of the criteria in each set (Table I). These findings (7,8) enabled the construction of a short questionnaire to screen for personal evidence that the symptom could have been caused by the blamed food and so to feed back advice on what to do about the symptom, tailored by the individual respondent's answers. One version of the resulting evidence-based self-care advice tool accompanies this article.

The first two questions within this advice tool are designed to identify those people who are recalling an actual episode when the symptom followed eating the food. In our community sample, 38% of interviewees (24% of food-symptom reports) would have been able to give a precise answer to both of these two questions for at least one food they blamed for a symptom. Such people should be advised to go to a hospital clinic for further investigation if the symptom and/or the food are important.

Those who cannot answer both of the first two questions precisely are asked to go on to questions 3a, 3b and 3c in the tool. Answering these helps the tool-user to consider other possible causes of the food-attributed symptom and whether the co-occurrence of food and symptom was mere coincidence. Among those of our interviewees who would have been referred to this part of the questionnaire, a further 24% (29% of food-symptom reports) had reason to believe that the co-occurrence of food and symptom was not a coincidence. These people should also be advised to ask their GP to refer them for further investigation by an appropriate specialist.

The reasoning offered after answering these questions identifies to the tool-users where they have a solid case against the food or where the attribution of the symptom to the food is likely to have been an understandable mistake. A very small proportion of those in the community who believe that they have a food allergy or sensitivity may have serious

emotional problems with eating what they perceive to be a harmful food. Such people are liable to find this tool no more persuasive than any other straight reasoning, let alone authoritative instruction. Where a powerful emotional reaction is discrete to the food, a clinical psychologist may be consulted about the possibility of therapy of the kind that is effective in reducing other sorts of phobia or panic. Contrary to the term “food aversion” (2), many of our interviewees showed no aversion to the food that they feared or professed to avoid; nevertheless, if a wholesome food has become disliked, similar “extinction” therapy may be applicable. Of course, where the disruptive emotion about a food is part of a more general problem about eating and perhaps about other parts of life, these wider difficulties should be addressed by the appropriate evidence-based therapeutic approach.

Copies of the accompanying tool may be distributed to members of the public by individual professionals, commercial or not-for-profit organisations or the media. However, the authors have retained copyright because the tool is valid only when used as a whole and so permission will not be given to reproduce a subset of the questions. Nevertheless, the layout of the material and the exact wording of the advice might be adjusted, so long as the authors review the edited version for any conflict with the psychological evidence. Requests to reproduce the tool also permit the authors to seek opportunities to assess its impact on individual consumers and on relevant aspects of the food supply and the health services.

ACKNOWLEDGEMENTS

This research project was funded by MAFF, London. The Principal Investigators were Professors D.A. Booth (Psychology) and I.W. Booth (Child Health), with Dietetics Consultant Mrs A. Macdonald. The Research Executives were Drs R.C. Knibb (Psychology) and A. Armstrong (Nutrition) and Mrs R.G. Platts. Dr Knibb is now Lecturer in Health Psychology, School of Nursing, University of Nottingham Medical Centre.

REFERENCES

1. Young, E., Stoneham, M.D., Petrukevitch, A., Barton, J., Rona, R.A. Population study of food intolerance. *Lancet*; 1994; 343; 1127-30.
2. Royal College of Physicians and British and British Nutrition Foundation. A Report on Food Intolerance and Food Aversion. *Journal of the Royal College of Physicians of London*; 1984; 18; No. 2; 83-123.
3. Knibb, R.C., Armstrong, A.M., Booth, D.A., Platts, R.G., Booth, I.W., Macdonald, A. Psychological characteristics of people with perceived food intolerance in a community sample. *Journal of Psychosomatic Research*; 1999; 47; 545-54.
4. Emsley, J., Fell, P. Was it something I ate? Food intolerance: what causes it and how to avoid it. Oxford University Press; 2000.
5. Booth, D.A. Evidence-based reduction of obesity: identification of a subculture's least fattening eating patterns. In: Workshop of IUNS Committee on Food Habits & Nutrition, International Congress of Nutrition, Montreal. *Appetite*; 1999; 32; 80-5.
6. Booth, D.A., Platts, R.G. Tool for assessing and reducing an individual's fat intake. *Appetite*; 2000; 34, 107-8.
7. Knibb, R., Booth, D.A., Armstrong, A., Platts, R., Macdonald, A., Booth, I.W. Episodic and semantic memory in reports of food intolerance. *Applied Cognitive Psychology*; 1999; 13; 451-64.
8. Booth, D.A., Knibb, R.C., Platts, R.G., Armstrong, A.M., Macdonald, A., Booth, I.W. The psychology of perceived food intolerance. *Leatherhead Food RA Food Industry Journal*; 1999; 2; No. 3; 325-34.

Table 1. Percentage of interviewees and reports categorised on a criterion used as a question in the tool, with percentages of those interviewees or reports scoring below the median for the total score on all criteria for actuality and for contingency of food-symptom events.

				Below median score for ACTUALITY		Below median score for CONTINGENCY	
		% of all interviewees		% of those people answering no			
Question number		people	reports	people	reports	people	reports
1	No	77	74	73	69	80	80
2	No	13	9	82	84	64	69
		% of those answering 'No' to Q1 or Q2		% of those answering 'No' to Q1/Q2 and 'Yes' to part or all of Q3			
3a	Yes	20	24	64	72	81	89
3b	Yes	27	31	73	78	85	93
3c	Yes	67	74	73	70	91	91
3a+b+c	Yes	7	7	58	63	92	97

Note – All interviewees (100%) who would have answered 'Yes' to Questions 3b or 3c had below the top score for Contiguity of eating the food to occurrence of the symptom.

Might you have a **food allergy** or a **sensitivity** to a food?

Millions of people blame foods for illnesses they get (or that their child has had). That is, they believe they have a **food intolerance**.

Do you (or your child) react badly after eating a food?
Do you suspect a food of causing a nasty symptom?

If so, you can sort out what's best to do, by answering the following queries.

The advice coming from your answers is based on scientific research with hundreds of volunteers who thought that something they ate or drank gave them an unpleasant symptom. (If it's a young child who may have a food allergy or sensitivity, then whoever most looks after the youngster can adapt these questions and act on the advice on the child's behalf.)

You may have more than one symptom that a food (or drink) may induce, or you may suspect more than one food. The following questions should be answered for just one symptom, and for just one food you believe is causing it. Go through the questions first for the food and the symptom that matter most.

Question 1

How many times has consuming the suspected food or drink been followed by the unpleasant symptom?

If you can give an *exact* number (for example, "Just once" or "Three times"), then go to Question 2 (at the top of the next page).

If you are *not* sure of the exact number of times it has happened (e.g., your answer is "a few times" or "regularly"), then skip the next page and go on to Question 3, on the third page.

Question 2

Can you remember whether or not you were ever able to consume the suspected food or drink and not get this symptom?

If you can't definitely remember *whether* there was such an occasion *or not*, go straight to Question 3 on the next page.

If you are quite clear
EITHER that you did *not* consume that food or drink on any occasion before this symptom first occurred
OR that you definitely *did* consume it before the occasion that was first followed by the symptom,
then this fits in with your precise answer to Question 1.

Your precision of answers to Questions 1 and 2 are evidence that, in the past as you recall it, the symptom followed the consumption of that food or drink on one or more occasions.

Such specific memories indicate that your belief that the food causes the symptom is based on one or more actual experiences. It is not just a guess – for example, from remembering the food or drink and remembering the symptom but without being sure that they ever went together.

With such definite answers to Questions 1 and 2, you do not have to answer any more questions. You have clear evidence that the foodstuff or beverage may be causing you problems. If having this symptom or avoiding that food or drink is bothersome, you should consider getting tested.

You should ask for tests for allergy or other sensitivity in a hospital in your Health region. You can phone or write direct, or you can ask your GP to refer you. Note that the tests given by other sorts of practitioners (e.g. hair analyses, bioelectric machines) are not scientifically founded and can give most misleading results.

Question 3

If you have followed the instruction to skip here after answering Question 1 or Question 2, it would be wise to think about the answers to the following parts A, B and C of this 3rd question before deciding what to do in future about the food that you suspect.

Question 3A

Do you sometimes get the symptom without consuming the food / drink?

If you do, isn't this evidence that the symptom can be caused by something else beside the food?

The nasty effect may come from another food, or it may have nothing to do with anything specific that you ate or drank.

If this symptom is bothersome, it would be wise to ask your GP for some advice other than avoiding that particular food or drink.

Question 3B

Do you sometimes consume the suspected food /drink without getting the symptom?

If that's so, isn't it evidence that this particular food or drink doesn't ever cause that symptom?

The unpleasant effect may arise from some other food. The suspected food or drink might be interacting with something else that could be tackled. The bad reaction might not from foods or drinks at all.

Consider asking your GP for advice about this symptom which doesn't involve that food or even the diet at all.

[Continued over the page]

Question 3C

Do you suspect two or three foods of causing the symptom, or perhaps quite a number of foods?

You know that you've eaten each of those foods and you've had the unpleasant symptom. Yet it's quite possible to be mis-led by sheer coincidence. Consider the fact that we usually eat something every few hours during the day. Therefore, a symptom that has nothing to do with any food can occur within some hours or even minutes of the foods in a meal or snack. So it is easy for someone very sensible to make a mistake about what is going on in such a complicated situation.

It would be advisable for you to consider the possibility that your symptom may come from only one of these foods or drinks. It may even have nothing to do with any of those particular items.

If the symptom is serious enough to take to a doctor, you should consider asking your GP for advice or treatment. If there is good reason from other cases to suspect something in the diet or in the environment, you can be referred or go direct to a hospital clinic for the testing of foods, pollens, dusts etc. that are known to provoke allergies or other sensitivities.

Do not trust the results of hair analyses, bio-electric machines or other scientifically unfounded tests: the conclusions can be badly misleading and cause you unnecessary trouble.

If you suspect **another food or drink** besides the one for which you answered the above questions, or if you have **another symptom** that you suspect a food of causing, you can go back to Question 1 and review your personal experience of that food and symptom.

(C) R.C. Knibb & D.A. Booth, University of Birmingham (UK), May 2000. To be reproduced as a whole only, with permission.

[If you or your doctor or other advisor has any general comment or question about this self-care advice tool, the Birmingham University prof. David Booth would be glad to receive a letter. Please note however that advice on individual circumstances cannot be provided.]