

## Dr Eugenio Sanchez-Moran

Lecturer

[School of Biosciences \(/schools/biosciences/index.aspx\)](/schools/biosciences/index.aspx)

### Contact details

**Telephone** +44 (0)121 41 45918 (tel:+44 121 41 45918)

**Fax** +44 (0)121 41 45925

**Email** [e.sanchezmoran@bham.ac.uk](mailto:e.sanchezmoran@bham.ac.uk) (mailto:e.sanchezmoran@bham.ac.uk)

School of Biosciences  
University of Birmingham  
Edgbaston  
Birmingham  
B15 2TT  
UK



### Qualifications

- 1996 B.A. Universidad Complutense de Madrid, Spain
- 2001 Ph.D. Universidad Complutense de Madrid, Spain
- 2002-2003 Marie-Curie Individual Fellowship, University of Birmingham, UK
- 2004-2008 BBSRC Postdoc Research Fellow, University of Birmingham, UK
- 2008 David Phillips Fellowship BBSRC, University of Birmingham, UK

### Postgraduate supervision

For a list of possible PhD projects offered by Dr Sanchez-Moran:

[www.findaphd.com/search/customlink.asp?inst=birm-Biol&supersurname=Sanchez-Moran](http://www.findaphd.com/search/customlink.asp?inst=birm-Biol&supersurname=Sanchez-Moran) (<http://www.findaphd.com/search/customlink.asp?inst=birm-Biol&supersurname=Sanchez-Moran>)

### Research

Research Theme: Molecular and Cell Biology

Short research description: Organisation and behaviour of chromosomes in plant meiosis

Full research description:

Our research aims to unravel the biological significance that the different levels of DNA compaction structures and components have on chromosome condensation and DNA processes in the nucleus. We believe that this research will contribute to the understanding of different important themes like cell division, cancer, stem cells, chromosome alterations, fertility and, plant and animal, breeding.

#### WHY DO WE DO IT?

The role of DNA is to store an individual's genetic information such that it can be used during normal growth and development and be accurately copied during the different divisions of the cell. Human cells contain DNA totalling about 2 m in length that has to be packed within the cell nucleus which is only 0.01 mm in diameter. Importantly, the DNA must be organised in such a way that it is readily accessible for a variety of crucial processes. The information it contains must be easily read (transcription) so that the cell can rapidly produce proteins. It must be readily duplicated (DNA replication) and accurately separated during cell division (mitosis) and sexual reproduction (meiosis). Also, it is essential that any break, knot or tangle that might occur can be repaired (DNA repair). DNA associates with different proteins forming a nucleo-protein complex called chromatin. This enables the compaction necessary to fit the naked DNA inside the cell nucleus whilst maintaining access to the genetic information. The chromatin is divided into individual structures constituting chromosomes. During the process of cell division when the individual chromosomes have been duplicated chromosome condensation is necessary to ensure their accurate distribution. Miss-regulation of chromosome condensation can lead to cell death, cancer and improper chromosome segregation during cell cycle or during the production of gametes.

#### WHAT DO WE KNOW ABOUT IT?

There are different levels of compaction involved in packaging DNA into chromosomes. The basic structure is the nucleosome, formed by wrapping naked DNA around a core of proteins known as histones. The nucleosomes are arranged along the DNA forming a 10nm diameter fibre, likened to beads on a string. Despite the old impression that nucleosomes were static structures, nowadays, a nucleosome is considered as a highly dynamic assemblage. Changes to this organised structure are facilitated through histone modifications, modelling factors and exchange of histone proteins. The nucleosomal fibre is further compacted by winding it into a 30 nm fibre whose structure remains controversial. This fibre is additionally arranged into loops that are attached to a multi-protein axis called the chromosome scaffold. Although the biochemistry of histones and other chromosome-associated proteins has been studied intensively, their interactions to achieve chromosome condensation are still poorly understood.

#### HOW DO WE DO IT?

The key proteins involved in chromosome condensation are conserved throughout eukaryotic evolution indicating that they are likely to have fundamental roles that are species-independent. We are using *Arabidopsis thaliana*, a plant model organism for basic research in genetics and molecular biology and a good experimental system without any of the ethical issues related to working with animals. We are also considering to start pilot experiments with other model systems to understand the evolution of chromatin structures. We are using a multidisciplinary approach combining new high-resolution cytogenetic techniques, mutant characterisation, proteomic analysis, and systems biology to resolve the complicated interactions of individual chromatin components that result in accurate chromosome condensation.

#### WHO IS FUNDING US?

## Publications

- Armstrong SJ, Sanchez-Moran E, Franklin FC. Cytological analysis of *Arabidopsis thaliana* meiotic chromosomes. *Methods in Molecular Biology* (2009) 558:131-45.
- Osman K, Sanchez-Moran E, Mann SC, Jones GH, Franklin FC. Replication protein A (AtRPA1a) is required for class I crossover formation but is dispensable for meiotic DNA break repair. *The EMBO Journal* (2009) 28(4):394-404.
- Sanchez-Moran E, Osman K, Higgins JD, Pradillo M, Cuáñado N, Jones GH, Franklin FCH. ASY1 co-ordinates early events in the plant meiotic recombination pathway. *Cytogenetic and Genome Research* (2008) 120: 302-312.
- Uanschou C, Siwiec T, Pedrosa-Harand A, Kerzendorfer C, Sanchez-Moran E, Novatchkova M, Akimcheva S, Klein F, Schläpfer P. A novel plant gene essential for meiosis is related to the human CtIP and the yeast COM1/SAE2 gene. *The EMBO Journal* (2007) 26: 5061-5070.
- Sanchez-Moran E, Santos JL, Jones GH, Franklin FCH. ASY1 mediates AtDMC1-dependent interhomolog recombination during meiosis in *Arabidopsis*. *Genes & Development* (2007) 21: 2220-2233.
- Pradillo M, Lopez E, Romero C, Sanchez-Moran E, Cuáñado N, Santos JL. An analysis of univalent segregation in meiotic mutants of *Arabidopsis thaliana*: A possible role for Synaptonemal Complex. *Genetics* (2007) 175: 505-511.
- Franklin FCH, Higgins JD, Sanchez-Moran E, Armstrong SJ, Osman K, Jackson N and Jones GH. Control of meiotic recombination in *Arabidopsis*: role of the MutL and MutS homologues. *Biochemical Society Transactions* (2006) 34: 542-544.
- Jackson N, Sanchez-Moran E, Buckling E, Armstrong SJ, Jones GH, Franklin FCH. Reduced meiotic crossovers and delayed prophase I progression in *AtMlh3*-deficient *Arabidopsis*. *The EMBO Journal* (2006) 25(6): 1315-1323.
- Osman K, Sanchez-Moran E, Higgins JD, Jones GH, Franklin FC. Chromosome synapsis in *Arabidopsis*: analysis of the transverse filament protein ZYP1 reveals novel functions for the synaptonemal complex. *Chromosoma* (2006) 115: 212-219.
- Higgins JD, Sanchez-Moran E, Armstrong SJ, Jones GH, Franklin FC. The *Arabidopsis* synaptonemal complex protein ZYP1 is required for chromosome synapsis and normal fidelity of crossing over. *Genes & Development* (2005) 19(20): 2488-500.
- Sanchez-Moran E, Higgins J, Mercier R, Armstrong SJ, Jones GH, Franklin FCH. Defining the meiotic proteome in *Brassica* and *Arabidopsis*. *Cytogenetic and Genome Research* (2005) 109(1-3): 181-9.
- Sanchez-Moran E, Jones GH, Franklin FCH, Santos JL. A Puromycin-Sensitive Aminopeptidase is essential for meiosis in *Arabidopsis thaliana*. *Plant Cell* (2004) 16: 2895-2909.
- Santos JL, Alfaro D, Sanchez-Moran E, Armstrong SJ, Franklin FCH, Jones GH. Partial diploidisation of meiosis in autotetraploid lines of *Arabidopsis thaliana*. *Genetics* (2003) 165: 1533-1540.
- Sanchez-Moran E, Armstrong SJ, Santos JL, Franklin FCH, Jones GH. Variation in chiasma frequency among eight accessions of *Arabidopsis thaliana*. *Genetics* (2002) 162: 1415-1422.
- Sanchez-Moran E, Armstrong SJ, Santos JL, Franklin FCH, Jones GH. Chiasma formation in *Arabidopsis thaliana* accession Wassileskija and in two meiotic mutants. *Chromosome Research* (2001) 9 (2): 121-128.
- Cuáñado N, Sanchez-Moran E, Barrios J, Santos JL. Searching for telomeric sequences in two *Allium* species. *Genome* (2001) 44 (4): 640-643.
- Sanchez-Moran E, Benavente E, Orellana J. Analysis of karyotypic stability of homoeologous-pairing (ph) mutants in allopolyploid wheats. *Chromosoma* (2001) 110 (5): 371-377.
- Sanchez-Moran E, Benavente E, Orellana J. Simultaneous identification of A, B, D and R genomes by genomic in situ hybridization in wheat-rye derivatives. *Heredity* (1999) 83 (3): 249-252.

