## PATIENT SAFETY INFORMATION CARD



Soliris can lower the ability of your immune system to fight infections, **especially meningococcal infection**, **which requires <u>immediate</u> medical attention**. If you experience any of the following symptoms, you should <u>immediately call your doctor</u>.

If you cannot reach your doctor, go to an Accident and Emergency department and show them this card.

- Headache with nausea or vomiting
- Headache with a stiff neck or stiff back
- Fever (raised temperature)
- Rash
- Confusion
- Muscle aches with flu-like symptoms
- Eyes sensitive to light



Seek emergency medical care immediately if you have <u>any</u> of these signs or symptoms and show this card.

**Even if you stop using Soliris,** keep this card with you for 3 months after your last Soliris dose. Your risk of meningococcal infection may continue for a long time after your last dose of Soliris.

## PATIENT SAFETY INFORMATION CARD



## Information for the Treating Doctor



This patient has been prescribed Soliris® (eculizumab) therapy, which increases the patient's susceptibility to meningococcal infection (Neisseria meningitidis) or other general infections.

- Meningococcal infections may become rapidly lifethreatening or fatal if not recognised and treated early
- Evaluate immediately if infection is suspected and treat with appropriate antibiotics if necessary
- Contact prescribing doctor (below) as soon as possible

For more information about Soliris, please refer to the full Summary of Product Characteristics or e-mail: medicalinformation.uk@alexion.com
In case of safety concerns, call 0800 321 3902 in UK, 1 800 936 544 in Ireland



## Patients receiving Soliris should carry this card at all times

Patient name	
Hospital where	treated
Doctor name	
Tel. number _	
Vaccination dat	•

