

Anti-anxiety drugs

Lorazepam (lor-az-eh-pam)

You should read this leaflet with your carer/caregiver.



Lorazepam:

- Is also called Ativan.
- Come as tablets.

- Lorazepam is used for anxiety, sleep problem and epilepsy.
- Lorazepam is sometimes used for people who are distressed. There is no clear evidence that it improves someone's behaviour.



The label on your lorazepam packet should tell you when and how much medicine to take.

Swallow your tablet with water.

If you forget to take your medicine, tell someone straight away.

You should carry on taking your medicine even if you feel better unless your doctor asks you to stop.



Drinking alcohol when taking lorazepam could make you feel sleepy.

Some medicines do not work well together.

You should tell your doctor about any other medicines you are taking.

Side effects:

- Medicine can have some effects on your body that are not wanted.
- Not everyone taking a medicine will get them.
- Many will go away with time.
- Many are rare.

You might want to talk to your doctor or the pharmacist or your key worker if you have any side effect.

These are some (but NOT all) side effects of lorazepam...



- Feeling unsteady on your feet
- Feeling sleepy and sluggish
- Feeling confused
- Problem with memory
- Feeling light headed and faint
- Feeling more aggressive
- Headache

Lorazepam should not be used if you have a lung disease, muscle weakness, liver disease, sleep disorder called sleep apnoea, drink alcohol excessively or have kidney problem.

Some side effects are more serious.

You should tell someone straight away if you have...

- Problems with breathing
- You see or hear things that are not there
- Rash or blotches on your skin



You should talk to your doctor before you stop taking your medicine.

Sometimes, if you stop taking your medicine, you may have:

- Shaking and sweating
- Anxiety or feel on edge
- Problem with sleeping
- Shaky hands
- Confusion or problem with memory
- Weight loss
- Stiffness and pain in muscles



Lorazepam may cause addiction. So, difficult to stop if used for long.

You may find the same dose of medicine is not working for you. **You should tell someone straight away if this happens to you.**



This leaflet does not tell you everything about lorazepam.

You can get more information from:

- your pharmacist, doctor, nurse or carer/ care giver
- check <http://www.choiceandmedication.org/cnwl/>
- NHS on 111
- medicine information helpline on 020 3317 5090.

This leaflet can be downloaded free of charge from <https://spectrom.wixsite.com/project/>

While preparing this leaflet we checked information in...

- The British National Formulary (www.bnf.org)
- Dulcan, M. K., & Ballard, R. (2015). *Helping parents and teachers understand medications for behavioural and emotional problems: A resource book of medication information handouts*. American Psychiatric Publishing, Washington DC, USA.

This leaflet is not designed for people with intellectual/learning disabilities to read on their own. Support (care) staff or care givers should go through the leaflet with the person who has intellectual disabilities.

All images copyright ©Photosymbols.